



The class of absolute decomposable inequality measures

Kristof Bosmans^{a,*}, Frank A. Cowell^b

^a Department of Economics, Maastricht University, Tongersestraat 53, 6211 LM Maastricht, The Netherlands

^b STICERD, London School of Economics, Houghton Street, London WC2A 2AE, United Kingdom

ARTICLE INFO

Article history:

Received 25 February 2009

Received in revised form 25 August 2010

Accepted 15 September 2010

Available online 24 September 2010

JEL classification:

D63

H20

H21

Keywords:

Inequality measures

Decomposability

Translation invariance

ABSTRACT

We provide a parsimonious axiomatisation of the complete class of absolute inequality indices. Our approach uses only a weak form of decomposability and does not require *a priori* that the measures be differentiable.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

Absolute inequality indices have the property that, for any income distribution, if any given number of dollars is added to (or subtracted from) every income, inequality remains unchanged. This means that such indices have the property that they are defined for negative incomes: this is particularly important in empirical applications where one is interested in applying inequality indices to the distribution of wealth (net worth may often be substantially negative) or to specific components of income (for example business income). Decomposability of inequality indices requires that, for an arbitrary subgroup of the population, if inequality in the subgroup increases then, *ceteris paribus*, inequality overall increases. Previous descriptions and characterizations of the class of indices that has these properties have invoked additional assumptions about structure – see Section 4 below. Our approach (in Sections 2 and 3) is deliberately minimalist in that it uses just the axioms required to define (a) an inequality measure (b) the “absolutist” property and (c) the population structure.

2. Setting

The income of individual i is a real number x_i , and the income distribution for a population of n individuals is a vector $\mathbf{x} := (x_1, x_2, \dots, x_n)$ in \mathbb{R}^n . The set of income distributions for m or more

individuals is $X_m := \cup_{n=m}^{\infty} \mathbb{R}^n$. For an income distribution \mathbf{x} , we write $n(\mathbf{x})$ for the dimension and $\mu(\mathbf{x})$ for the mean. A vector of dimension n of which all components are equal to 1 is denoted by $\mathbf{1}_n$.

An *inequality measure* is a continuous function $I: X_1 \rightarrow \mathbb{R}$ with the property that $I(\mathbf{x}) = 0$ if \mathbf{x} is an equal income distribution. The higher the value of the function I , the higher income inequality. To give meaning to I it is standard to assume the following two properties:

Anonymity. For every $\mathbf{x} \in X_1$, $I(\mathbf{x}) = I(\mathbf{x}')$ if \mathbf{x}' is obtained from \mathbf{x} by rearrangement of components.

Transfer principle. For every $\mathbf{x} \in X_2$ and any positive real number δ , we have that if $x_i < x_i + \delta \leq x_j - \delta < x_j$, then $I(x_1, \dots, x_i, \dots, x_j, \dots, x_n) > I(x_1, \dots, x_i + \delta, \dots, x_j - \delta, \dots, x_n)$.

The essential property for an *absolute* inequality measure is this:

Translation invariance. For every $\mathbf{x} \in X_1$ and any real number δ ,

$$I(\mathbf{x}) = I(\mathbf{x} + \delta \mathbf{1}_n). \quad (1)$$

Finally, the following two axioms describe the relationship between population structure and inequality:

Decomposability. There exists a function A such that, for all $\mathbf{x}, \mathbf{y} \in X_2$,

$$I(\mathbf{x}, \mathbf{y}) = A(I(\mathbf{x}), I(\mathbf{y}), \mu(\mathbf{x}), \mu(\mathbf{y}), n(\mathbf{x}), n(\mathbf{y})),$$

where A is continuous and strictly increasing in its first two arguments.

Replication invariance. For every $\mathbf{x} \in X_1$, $I(\mathbf{x}, \mathbf{x}) = I(\mathbf{x})$.

* Corresponding author. Tel.: +31 43 3883942; fax: +31 43 3884878.

E-mail addresses: k.bosmans@maastrichtuniversity.nl (K. Bosmans), f.cowell@lse.ac.uk (F.A. Cowell).

3. Result

Our result characterizes the class of inequality measures satisfying the five properties listed in Section 2.

Theorem. *An inequality measure I satisfies anonymity, the transfer principle, replication invariance, decomposability and translation invariance if and only if there exists a real number c and a continuous and strictly increasing function $f: \mathbb{R} \rightarrow \mathbb{R}$, with $f(0) = 0$, such that, for all $\mathbf{x} \in X_1$,*

$$f(I(\mathbf{x})) = \begin{cases} \frac{1}{n(\mathbf{x})} \sum_{i=1}^{n(\mathbf{x})} \{e^{c[x_i - \mu(\mathbf{x})]} - 1\} & \text{if } c \neq 0, \\ \frac{1}{n(\mathbf{x})} \sum_{i=1}^{n(\mathbf{x})} [x_i - \mu(\mathbf{x})]^2 & \text{if } c = 0. \end{cases} \quad (2)$$

Proof. The inequality measures given in Eq. (2) satisfy anonymity, the transfer principle, replication invariance, decomposability and translation invariance. We focus on the reverse implication.

If I satisfies anonymity, the transfer principle, replication invariance and decomposability, then (Shorrocks, 1984, Theorem 4) there exist functions F and ϕ such that

$$F(I(\mathbf{x}), \mu(\mathbf{x})) = \frac{1}{n(\mathbf{x})} \sum_{i=1}^{n(\mathbf{x})} [\phi(x_i) - \phi(\mu(\mathbf{x}))], \quad (3)$$

where ϕ is continuous and strictly convex, and where F is continuous in I and μ , strictly increasing in I , with $F(0, \mu) = 0$. Define $\hat{F}(I, \mu) := nF(I, \mu)$. Choose any \mathbf{x}_1 and \mathbf{x}_2 such that $n(\mathbf{x}_1) = n(\mathbf{x}_2) = 2$, and write $\mathbf{x}_{12} := (\mathbf{x}_1, \mathbf{x}_2)$, $I_k := I(\mathbf{x}_k)$ and $\mu_k := \mu(\mathbf{x}_k)$. We may write

$$I(\mathbf{x}_{12}) = H(\hat{F}(I_1, \mu_1) + \hat{F}(I_2, \mu_2), \mu_1, \mu_2), \quad (4)$$

where H is a function implicitly defined by

$$\hat{F}(H(z, \mu_1, \mu_2), \mu_{12}) = z + 2\phi(\mu_1) + 2\phi(\mu_2) - 4\phi(\mu_{12}).$$

Define $\lambda := e^\delta$ and $\theta_k := e^{\mu_k}$. Then $\lambda\theta_k = e^{\mu_k + \delta}$, and Eqs. (4) and (1) imply

$$\begin{aligned} & H(\hat{F}(I_1, \ln(\theta_1)) + \hat{F}(I_2, \ln(\theta_2)), \ln(\theta_1), \ln(\theta_2)) \\ &= H(\hat{F}(I_1, \ln(\lambda\theta_1)) + \hat{F}(I_2, \ln(\lambda\theta_2)), \ln(\lambda\theta_1), \ln(\lambda\theta_2)), \end{aligned}$$

or, with appropriate definitions of G from \hat{F} and J from H ,

$$J(G(I_1, \theta_1) + G(I_2, \theta_2), \theta_1, \theta_2) = J(G(I_1, \lambda\theta_1) + G(I_2, \lambda\theta_2), \lambda\theta_1, \lambda\theta_2).$$

From Lemma 1 in Shorrocks (1984):

$$\begin{aligned} G(I_1, \lambda\theta_1) &= cG(I_1, \theta_1), \\ G(I_2, \lambda\theta_2) &= cG(I_2, \theta_2), \end{aligned}$$

from which it follows that

$$c = \frac{G(I, \lambda\theta)}{G(I, \theta)} = c(\lambda),$$

independent of I and θ . Therefore

$$\begin{aligned} G(I, \lambda\theta) &= c(\lambda)G(I, \theta) \\ &= c(\lambda)c(\theta)G(I, 1) \\ &= c(\lambda\theta)G(I, 1). \end{aligned}$$

Because $c(\lambda\theta) = c(\lambda)c(\theta)$ for all positive λ and θ , we have $c(\theta) = \theta^c$ (Aczél, 2006, Theorem 4, p. 144). Consequently

$$\begin{aligned} G(I, \theta) &= \theta^c G(I, 1), \\ \hat{F}(I, \mu) &= e^{c\mu} \hat{F}(I, 0), \end{aligned}$$

and Eq. (3) yields

$$\hat{F}(I(\mathbf{x}), 0) = e^{-c\mu} \sum_{i=1}^{n(\mathbf{x})} [\phi(x_i) - \phi(\mu)]. \quad (5)$$

In the case of two individuals,

$$0 = \hat{F}(I(\mathbf{x} + \delta \mathbf{1}_2), \mu(\mathbf{x}) + \delta) - e^{c\delta} \hat{F}(I(\mathbf{x}), \mu(\mathbf{x})),$$

and writing $\mathbf{x} := (u, v)$ this becomes

$$\begin{aligned} 0 &= \left[\phi(u + \delta) + \phi(v + \delta) - 2\phi\left(\frac{u + v}{2} + \delta\right) \right] \\ &\quad - e^{c\delta} \left[\phi(u) + \phi(v) - 2\phi\left(\frac{u + v}{2}\right) \right]. \end{aligned} \quad (6)$$

If we define

$$\psi(u, \delta) := \phi(u + \delta) - e^{c\delta} \phi(u), \quad (7)$$

then Eq. (6) becomes

$$0 = \psi(u, \delta) + \psi(v, \delta) - 2\psi\left(\frac{u + v}{2}, \delta\right).$$

This is a Pexider equation with solution (Aczél, 2006, p. 142)

$$\psi(u, \delta) = a(\delta)u + b(\delta). \quad (8)$$

From Eq. (7),

$$\begin{aligned} \psi(u, \mu + \delta) &= \phi(u + \mu + \delta) - e^{c(\mu + \delta)} \phi(u) \\ &= \psi(u + \mu, \delta) + e^{c\delta} \phi(u + \mu) - e^{c(\mu + \delta)} \phi(u) \\ &= \psi(u + \mu, \delta) + e^{c\delta} \psi(u, \mu). \end{aligned} \quad (9)$$

Substitute from Eq. (8) into Eq. (9) to get

$$a(\mu + \delta)u + b(\mu + \delta) = a(\delta)(u + \mu) + b(\delta) + e^{c\delta}[a(\mu)u + b(\mu)],$$

which implies

$$a(\mu + \delta) = a(\delta) + e^{c\delta} a(\mu) = a(\mu) + e^{c\mu} a(\delta), \quad (10)$$

$$b(\mu + \delta) = a(\delta)\mu + b(\delta) + e^{c\delta} b(\mu) = a(\mu)\delta + b(\mu) + e^{c\mu} b(\delta). \quad (11)$$

There are two cases to consider.

Case 1. ($c \neq 0$). In this case Eq. (10) gives

$$a(\delta)(e^{c\mu} - 1) = a(\mu)(e^{c\delta} - 1),$$

$$a(\mu) = \frac{a(\delta)}{e^{c\delta} - 1} (e^{c\mu} - 1),$$

so that, for any given value of δ and all μ ,

$$a(\mu) = \frac{k_1}{k_2} (e^{c\mu} - 1),$$

where $k_1 := a(\delta)$ and $k_2 := e^{c\delta} - 1$ can be taken as constants (conditional on the chosen δ and the arbitrary value of c). So the only solution is

$$a(\mu) = \alpha (e^{c\mu} - 1).$$

Therefore, from Eq. (11)

$$b(\mu + \delta) = \mu\alpha(e^{c\delta} - 1) + b(\delta) + e^{c\delta}b(\mu) = \delta\alpha(e^{c\mu} - 1) + b(\mu) + e^{c\mu}b(\delta),$$

so that

$$\mu\alpha(e^{c\delta} - 1) + b(\mu)(e^{c\delta} - 1) = \delta\alpha(e^{c\mu} - 1) + b(\delta)(e^{c\mu} - 1),$$

$$b(\mu)(e^{c\delta} - 1) = [\delta\alpha + b(\delta)](e^{c\mu} - 1) - \mu\alpha(e^{c\delta} - 1),$$

$$b(\mu) = \frac{\delta\alpha + b(\delta)}{e^{c\delta} - 1}(e^{c\mu} - 1) - \mu\alpha.$$

Since this must be true for arbitrary δ and c , the solution is

$$b(\mu) = \beta(e^{c\mu} - 1) - \alpha\mu. \quad (12)$$

Case 2. ($c = 0$). Here Eq. (10) becomes

$$a(\mu + \delta) = a(\mu) + a(\delta),$$

and the solution to this Cauchy equation is $a(\mu) = \kappa\mu$ (Aczél, 2006, Theorem 2, pp. 35–36). Therefore, from Eq. (11)

$$b(\mu + \delta) = \kappa\mu\delta + b(\delta) + b(\mu). \quad (13)$$

Noting that $b(0) = 0$ and letting $\delta = -\mu$ and $\kappa = 2\gamma$, we get

$$b(\mu) + b(-\mu) = 2\gamma\mu^2,$$

to which the solution is

$$b(\mu) = \gamma\mu^2 + p(\mu),$$

where $p(\mu) := \Pi(\mu, -\mu)$ with the property $p(\mu) = -p(-\mu)$ (Polyanin and Manzhirrov, 1998). Plugging this into Eq. (13) gives

$$p(\mu + \delta) = p(\mu) + p(\delta),$$

to which the solution is $p(\mu) = \eta\mu$. Therefore

$$b(\mu) = \gamma\mu^2 + \eta\mu. \quad (14)$$

From Eq. (8), we have $\psi(0, x) = b(x)$; so from Eqs. (12) and (14) we find

$$\psi(0, x) = \begin{cases} \beta(e^{cx} - 1) - \alpha x & \text{if } c \neq 0, \\ \gamma x^2 + \eta x & \text{if } c = 0. \end{cases} \quad (15)$$

Note that, from Eq. (7), we have:

$$\phi(x) = \psi(0, x) + e^{cx}\phi(0). \quad (16)$$

However the constant $\phi(0)$ is arbitrary. Setting $\phi(0) = 0$ and substituting from Eqs. (15) and (16) into Eq. (5) produces Eq. (2). \square

4. Concluding remarks

While it is well known that the measures in Eq. (2) belong to the class of absolute decomposable measures (Chakravarty and Tyagarupananda, 1998, 2009), it has not previously been demonstrated that *only* these measures belong to the class: previous treatments have introduced *a priori* additional restrictions such as differentiability and have used a stronger decomposability property.

Letting $c = 0$, we obtain the variance and, letting $c < 0$, the family in Eq. (2) is ordinarily equivalent to the Kolm (1976) family of measures. As the parameter c decreases, the corresponding measures are more sensitive to transfers between incomes at the bottom of the income distribution. Note that the Kolm measures are the only members of the class of absolute decomposable measures satisfying *transfer sensitivity* (Shorrocks and Foster, 1987), which is the requirement that a rich-to-poor transfer combined with a simultaneous poor-to-rich transfer at a higher income level (such that there is no effect on the variance) decreases inequality.

Acknowledgment

We thank an anonymous referee for useful suggestions.

References

- Aczél, J., 2006. Lectures on Functional Equations and Their Applications. Dover Publications, Inc., Mineola.
- Chakravarty, S.R., Tyagarupananda, S., 1998. The subgroup decomposable absolute indices of inequality. In: Chakravarty, S.R., Coondoo, D., Mukherjee, R. (Eds.), Quantitative Economics: Theory and Practice, Essays in Honor of Professor N. Bhattacharya. Allied Publishers Limited, New Delhi, pp. 247–257.
- Chakravarty, S.R., Tyagarupananda, S., 2009. The subgroup decomposable intermediate indices of inequality. Spanish Economic Review 11, 63–93.
- Kolm, S.C., 1976. Unequal inequalities I. Journal of Economic Theory 12, 416–442.
- Polyanin, A.D., Manzhirrov, A.V., 1998. Handbook of Integral Equations: Exact Solutions. Faktorial, Moscow.
- Shorrocks, A.F., 1984. Inequality decomposition by population subgroups. Econometrica 52, 1369–1385.
- Shorrocks, A.F., Foster, J.E., 1987. Transfer sensitive inequality measures. Review of Economic Studies 54, 485–497.